

XL1-Blue Subcloning-Grade Competent Cells

Catalog #200130



MATERIALS PROVIDED

Materials provided	Quantity	Efficiency (cfu/μg of pUC18 DNA)
XL1-Blue subcloning-grade competent cells (blue tubes)	8 × 0.5 ml	≥1 × 10 ⁶
pUC18 control plasmid (0.1 ng/μl in TE buffer)	10 μl	—

Storage: Competent cells must be placed immediately at the bottom of a –80°C freezer directly from the dry ice shipping container. Do not store the cells in liquid nitrogen.

QUALITY CONTROL TESTING

Transformations are performed both with and without plasmid DNA using 50-μl aliquots of cells and 100 pg of pUC18 control plasmid following the protocol outlined below. Following transformation, 100-μl samples of the culture are plated in duplicate on LB agar plates with 100 μg/ml of ampicillin. The plates are incubated at 37°C overnight and the efficiency is calculated based on the average number of colonies per plate.

BACKGROUND

The XL1-Blue strain allows blue-white color screening for recombinant plasmids and is an excellent host strain for routine cloning applications using plasmid or lambda vectors.

XL1-Blue Genotype: *recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac* [F' *proAB lacI^qΔM15 Tn10* (Tet^r)]. (Genes listed signify mutant alleles. Genes on the F' episome, however, are wild-type unless indicated otherwise).

XL1-Blue cells are resistant to tetracycline. XL1-Blue cells are endonuclease (*endA*) deficient, which greatly improves the quality of miniprep DNA, and are recombination (*recA*) deficient, improving insert stability. The *hsdR* mutation prevents the cleavage of cloned DNA by the *EcoK* endonuclease system. The *lacI^qΔM15* gene on the F' episome allows blue-white color screening.

TRANSFORMATION PROTOCOL

1. Pre-chill two 14-ml BD Falcon polypropylene round-bottom tubes on ice. One tube is for the experimental transformation and one tube is for the pUC18 control. Preheat SOC medium[§] to 42°C.
2. Thaw the competent cells on ice. When thawed, gently mix and aliquot 50 μl of cells into each of the two pre-chilled tubes.
3. Add 0.1–50 ng of the experimental DNA (see *Quantity and Volume of DNA*, reverse page, for guidelines) to one aliquot of cells and add 1 μl of the pUC18 control DNA to the other aliquot. Swirl the tubes gently.
4. Incubate the tubes on ice for 20 minutes.
5. Heat-pulse the tubes in a 42°C water bath for 45 seconds. The duration of the heat pulse is **critical** for maximum efficiency.
6. Incubate the tubes on ice for 2 minutes.
7. Add 0.9 ml of preheated SOC medium and incubate the tubes at 37°C for 30 minutes with shaking at 225–250 rpm.
8. Plate ≤200 μl of the transformation mixture on LB agar plates containing the appropriate antibiotic (and containing IPTG and X-gal if color screening is desired).[§] For the pUC18 control transformation, plate 100 μl of the transformation mixture LB–ampicillin agar plates.[§]

Notes Cells may be concentrated by centrifuging at 1000 rpm for 10 minutes. Resuspend the pellet in 200 μl of SOC medium.

If plating <100 μl of cells, pipet the cells into a 200 μl pool of SOC medium and then spread the mixture with a sterile spreader. If plating ≥100 μl, the cells can be spread on the plates directly. Tilt and tap the spreader to remove the last drop of cells.

Some β-galactosidase fusion proteins are toxic to the host bacteria. If an insert is suspected to be toxic, plate the cells on media without X-gal and IPTG. Color screening will be eliminated, but lower levels of the potentially toxic protein will be expressed.

9. Incubate the plates at 37°C overnight (at least 17 hours for blue-white color screening). Colonies containing plasmids with inserts will remain white, while colonies containing plasmids without inserts will be blue. The blue color can be enhanced by incubating the plates for two hours at 4°C following the overnight incubation at 37°C.
10. For the pUC18 control, expect 10 colonies (≥1 × 10⁶ cfu/μg pUC18 DNA). For the experimental DNA, the number of colonies will vary according to the size and form of the transforming DNA, with larger and non-supercoiled DNA producing fewer colonies.

[§]See *Preparation of Media and Reagents*.

TRANSFORMATION GUIDELINES AND TROUBLESHOOTING

Storage Conditions: Competent and supercompetent cells are very sensitive to even small variations in temperature and must be stored at the bottom of a -80°C freezer. Transferring tubes from one freezer to another may result in a loss of efficiency.

Use of 14-ml BD Falcon polypropylene round-bottom tubes: It is important that 14-ml BD Falcon polypropylene round-bottom tubes (BD Biosciences Catalog #352059) are used for the transformation protocol, since other tubes may be degraded by β -mercaptoethanol. In addition, the duration of the heat-pulse is critical and has been optimized for these tubes.

Aliquoting Cells: Keep the cells on ice at all times during aliquoting. It is essential that the polypropylene tubes are placed on ice before the cells are thawed and that the cells are aliquoted directly into pre-chilled tubes. It is also important to use the volume of cells indicated in step 2 of the *Transformation Protocol*. Decreasing the volume will result in lower efficiencies.

Quantity and Volume of DNA: The greatest efficiency is obtained from the transformation of 1 μ l of 0.1 ng/ μ l supercoiled pUC18 DNA per reaction. A greater number of colonies may be obtained by transforming up to 50 ng DNA, although the resulting efficiency (cfu/ μ g DNA) may be lower. The volume of the DNA solution added to the reaction may be increased to up to 10% of the reaction volume, but the transformation efficiency may be reduced.

Heat Pulse Duration: Optimal transformation efficiency is observed when cells are heat-pulsed at 42°C for 45–50 seconds. Efficiency decreases sharply when cells are heat-pulsed for <45 seconds or for >60 seconds.

Blue-White Color Screening: Blue-white color screening for recombinant plasmids is available when transforming host strains that contain the *lacI^qΔM15* gene on the F' episome with a plasmid that provides α -complementation (e.g. Stratagene's pBluescript® II vector). When performing blue-white color screening, incubate the LB agar plates containing IPTG and X-gal at 37°C for at least 17 hours to allow color development. The blue color can be enhanced by subsequent incubation of the plates for two hours at 4°C.

PREPARATION OF MEDIA AND REAGENTS

SOB Medium (per Liter) 20.0 g of tryptone 5.0 g of yeast extract 0.5 g of NaCl Add deionized H ₂ O to a final volume of 1 liter Autoclave Add 10 ml of filter-sterilized 1 M MgCl ₂ and 10 ml of filter-sterilized 1 M MgSO ₄ prior to use	SOC Medium (per 100 ml) Note <i>This medium should be prepared immediately before use.</i> 2 ml of filter-sterilized 20% (w/v) glucose or 1 ml of filter-sterilized 2 M glucose SOB medium (autoclaved) to a final volume of 100 ml
LB Agar (per Liter) 10 g of NaCl 10 g of tryptone 5 g of yeast extract 20 g of agar Add deionized H ₂ O to a final volume of 1 liter Adjust pH to 7.0 with 5 N NaOH Autoclave	LB-Ampicillin Agar (per Liter) 1 liter of LB agar, autoclaved Cool to 55°C Add 10 ml of 10 mg/ml filter-sterilized ampicillin Pour into petri dishes (~25 ml/100-mm plate)

Preparation of Agar Plates for Blue-White Color Screening

To prepare plates for blue-white screening, prepare LB agar as indicated above. When adding the antibiotic, also add 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-gal) to a final concentration of 80 μ g/ml [prepared in dimethylformamide (DMF)] and isopropyl-1-thio- β -D-galactopyranoside (IPTG) to a final concentration of 20 mM (prepared in sterile dH₂O). Alternatively, 100 μ l of 10 mM IPTG and 100 μ l of 2% X-gal may be spread on solidified LB agar plates 30 minutes prior to plating the transformations. (For consistent color development across the plate, pipet the X-gal and the IPTG into a 100- μ l pool of SOC medium and then spread the mixture across the plate. Do not mix the IPTG and the X-gal before pipetting them into the pool of SOC medium because these chemicals may precipitate.)

LIMITED PRODUCT WARRANTY

This warranty limits our liability to replacement of this product. No other warranties of any kind, express or implied, including without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Stratagene. Stratagene shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product.

ENDNOTES

pBluescript® is a registered trademark of Stratagene in the United States.

Stratagene Technical Services

USA/Canada (Toll-free) 800 894 1304

Europe (Toll-free) 00800 7400 7400

Email tech_services@stratagene.com

World Wide Web www.stratagene.com