

AffinityScript QPCR cDNA Synthesis Kit

INSTRUCTION MANUAL

Catalog #600559

Revision C.01

For In Vitro Use Only

600559-12

LIMITED PRODUCT WARRANTY

This warranty limits our liability to replacement of this product. No other warranties of any kind, express or implied, including without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Agilent. Agilent shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product.

ORDERING INFORMATION AND TECHNICAL SERVICES

United States and Canada

Agilent Technologies
Stratagene Products Division

11011 North Torrey Pines Road
La Jolla, CA 92037

Telephone (858) 373-6300
Order Toll Free (800) 424-5444
Technical Services (800) 894-1304
Internet techservices@agilent.com
World Wide Web www.stratagene.com

Europe

Location	Telephone	Fax	Technical Services
Austria	0800 292 499	0800 292 496	0800 292 498
Belgium	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 15775	0800 15740	0800 15720
France	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 919 288	0800 919 287	0800 919 289
Germany	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 182 8232	0800 182 8231	0800 182 8234
Netherlands	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 023 0446	+31 (0)20 312 5700	0800 023 0448
Switzerland	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 563 080	0800 563 082	0800 563 081
United Kingdom	00800 7000 7000	00800 7001 7001	00800 7400 7400
	0800 917 3282	0800 917 3283	0800 917 3281

All Other Countries

Please contact your local distributor. A complete list of distributors is available at www.stratagene.com.

AffinityScript QPCR cDNA Synthesis Kit

CONTENTS

Materials Provided	1
Storage Conditions	1
Introduction	2
Kit Components.....	2
Options for QPCR Analysis	3
PreProtocol Considerations	4
RNA Isolation.....	4
Using a Reference RNA for QRT-PCR Experiments	4
cDNA Synthesis Reaction	5
The Reverse Transcriptase-Mediated Polymerase Chain Reaction Protocol	6
Synthesis of First-Strand cDNA Using Reverse Transcriptase	6
Real-Time PCR Quantification of First-Strand cDNA.....	7
Troubleshooting	8
References	9
Endnotes	9
MSDS Information	9

AffinityScript QPCR cDNA Synthesis Kit

MATERIALS PROVIDED

Materials provided	Concentration	Quantity ^a
cDNA Synthesis Master Mix	2×	500 µl
AffinityScript RT/ RNase Block Enzyme Mixture	—	50 µl
Oligo(dT) primer	100 ng/µl	15 µg
Random primers	100 ng/µl	15 µg
RNase-free H ₂ O	—	1.2 ml

^a The AffinityScript QPCR cDNA synthesis kit provides enough reagents for fifty 20-µl reactions.

STORAGE CONDITIONS

All Reagents: -20°C

Revision C.01

© Agilent Technologies, Inc. 2009.

INTRODUCTION

The AffinityScript QPCR cDNA Synthesis Kit is designed for the highest efficiency conversion of RNA to cDNA and is fully optimized for two-step quantitative reverse transcription-PCR (QRT-PCR) applications. A two-step RT-PCR format is useful for amplifying multiple targets from a single cDNA source, for maintaining archival cDNA, and for providing maximum flexibility in selecting a downstream QPCR reagent system. The AffinityScript QPCR cDNA synthesis kit provides a streamlined master mix format and fast protocol, with a 15-minute cDNA synthesis step and an overall protocol time of just 25 minutes. The kit is formatted for high-yield production of cDNA up to 12 kb and each reaction accommodates a range of RNA amounts from fg to µg.

Advantages of the AffinityScript QPCR cDNA Synthesis Kit

- Robust cDNA synthesis across a broad range of temperatures from 37–55°C
- High affinity for primer-template complexes promotes optimal cDNA yield
- Produces excellent yields of amplifiable cDNA
- Can synthesize long cDNA targets, up to 12 kb, from small amounts of input RNA
- Able to reverse transcribe through RNA secondary structures
- Strict enzyme purity specifications ensure the absence of contaminating RNase and exonuclease activities

Kit Components

The AffinityScript QPCR cDNA synthesis kit provides a comprehensive set of reagents necessary to generate high-quality, QPCR-ready cDNA templates from either poly(A)⁺ mRNA or total RNA. Key to the performance of the kit is the QPCR-grade AffinityScript Reverse Transcriptase (RT) enzyme and the corresponding QPCR-optimized RT buffer. The kit is formatted as a convenient master mix with few pipetting steps, saving you time and ensuring experiment-to-experiment reproducibility.

cDNA Synthesis Master Mix

The cDNA synthesis master mix contains a buffer that is specifically optimized for QRT-PCR performance, allowing a fast protocol and reducing variability in Ct measurements between reactions. In addition to the optimized buffer, the master mix contains MgCl₂ and dNTPs.

cDNA Primers

The cDNA priming strategy can affect cDNA yield, sensitivity, and detection of certain targets, such as GC-rich targets or sequences located at the 5' or 3' end of a transcript.¹ For this reason, individually packaged random nonamers and oligo(dT) primers are provided separately from the master mix, allowing you to use the best priming strategy for your specific target.²

AffinityScript RT/RNase Block Enzyme Mixture

The kit includes our QPCR-grade AffinityScript Multiple Temperature Reverse Transcriptase, a genetically engineered version of Moloney Murine Leukemia Virus Reverse Transcriptase (MMLV-RT) which is highly thermostable, allowing you to reverse transcribe at your preferred reaction temperature. AffinityScript RT is provided, in combination with RNase block, in a separate tube so that *no-RT control* reactions may be included in the QRT-PCR experiments. The RNase block serves as a safeguard against contaminating RNases.

QPCR-grade AffinityScript RT is stringently quality-controlled to verify the absence of nuclease contaminants that adversely affect cDNA synthesis, particularly from small input RNAs. It has been tested in QRT-PCR experiments to ensure sensitive and reproducible performance over a broad range of RNA template amounts and over a variety of RNA targets that vary in size, abundance, and GC-content.

Options for QPCR Analysis

The AffinityScript QPCR cDNA synthesis kit is designed for maximum flexibility in downstream QPCR analysis options, and is qualified for use with SYBR® Green-based and probe-based detection. The kit is tested in 2-step QRT-PCR using Stratagene QPCR master mix kits. See Table 1 for a list of the QPCR master mix kits available. Brilliant II QPCR kits are recommended for maximum sensitivity, while Brilliant II Fast QPCR kits are recommended for maximum speed.

TABLE 1 Recommended QPCR Analysis Kits

Detection Method	Application	QPCR Master Mix Kit	Stratagene Catalog #
SYBR® Green dye	Maximum sensitivity QPCR	Brilliant II SYBR® Green QPCR master mix	600828
	Rapid run-time QPCR	Brilliant II Fast SYBR® Green QPCR master mix	600843
Probe	Maximum sensitivity QPCR	Brilliant II QPCR master mix	600804
	Rapid run-time QPCR	Brilliant II Fast QPCR master mix	600845

PREPROTOCOL CONSIDERATIONS

RNA Isolation

High-quality intact RNA is essential for successful synthesis of full-length cDNA. Total RNA can be rapidly isolated and purified from cells using Stratagene Absolutely RNA isolation kits, which are available for nano-, micro- and miniprep scale RNA purifications (Catalog #400753, #400805, and #400800, respectively). Total RNA may also be isolated from formalin-fixed, paraffin-embedded (FFPE) tissues using the Absolutely RNA FFPE kit (Catalog #400809 and #400811).

Oligo(dT)-selection for poly(A)⁺ RNA is typically not necessary, although including this step may improve the yield of specific cDNA templates. The Absolutely mRNA purification kit (Catalog #400806) is recommended for this application.

For QRT-PCR experiments with mammalian cells, we offer the SideStep Lysis and Stabilization Buffer (Catalog #400900), which allows you to skip RNA isolation steps and analyze gene expression in cell lysates directly. Since the SideStep lysates also contain genomic DNA, it is critical to design PCR primers to span adjacent exons in order to selectively amplify cDNA and prevent amplification of the intron-containing genomic DNA.

Preventing RNase Contamination

Take precautions to minimize the potential for contamination by ribonucleases (RNases). RNA isolation should be performed under RNase-free conditions. Wear gloves and use sterile tubes, pipet tips, and RNase-free water. The RNase block that is included in the cDNA synthesis reaction mixture provides additional protection against RNase contamination.

Preventing Genomic DNA Contamination

Contaminating DNA can be removed from the RNA preparation using an RNase-free DNase. All of the Stratagene Absolutely RNA kits include RNase-free DNase which is used in a rapid on-column DNase treatment protocol. Additionally, PCR primers may be designed to span adjacent exons in order to prevent amplification of the intron-containing genomic DNA.

Using a Reference RNA for QRT-PCR Experiments

In order to reliably compare data across multiple experiments and instruments, it is essential to have a constant reference material to assess the performance of each QPCR run and to quantify gene expression levels. Stratagene QPCR Reference Total RNA is a high-quality control for quantitative PCR gene-expression analysis. Including a standard curve with the Stratagene QPCR Reference Total RNA in every experiment allows you to assess QPCR assay efficiency and precision, and to quantitate relative to an unchanging reference standard. In addition, the broad gene coverage allows you to use the reference material for nearly any human or mouse gene being investigated, thus eliminating the extra work required in generating new standards for each new gene target.

We offer QPCR Reference Total RNAs for both the human system (Catalog #750500) and the mouse system (Catalog #750600). The QPCR Human Reference Total RNA is composed of total RNA from 10 human cell lines with quantities of RNA from the individual cell lines optimized to maximize representation of gene transcripts present in low, medium, and high abundance. The QPCR Mouse Reference Total RNA is derived from RNA pooled from 11 mouse cell lines. These reference RNAs are carefully screened for contaminating genomic DNA, the presence of which can complicate interpretation of QRT-PCR assay data.

cDNA Synthesis Reaction

Primer Selection

The optimum primer type [oligo(dT) or random primer] varies for different targets and should be determined empirically with each target. Stratagene QPCR Reference Total RNA (available separately) can be used for this step. (See *Using a Reference RNA for QRT-PCR Experiments* for more details.) For most targets, the best results are achieved using either oligo(dT) or random primers. For some challenging targets (long or secondary structure-rich targets), however, using a mixture of oligo(dT) and random primers may increase cDNA yield. When testing the use of mixed primers, adding a mixture of 170 ng oligo(dT) primer and 30 ng random primers to each 20- μ l reaction is a good starting point.

Performing No-RT Control Reactions

Perform no-RT control reactions for each RNA sample by omitting AffinityScript RT/ RNase block from the reaction. The no-RT control is expected to generate no signal in subsequent QPCR if there is no amplification of genomic DNA. See *Preventing Genomic DNA Contamination in RNA Isolation*.

THE REVERSE TRANSCRIPTASE-MEDIATED POLYMERASE CHAIN REACTION PROTOCOL

Synthesis of First-Strand cDNA Using Reverse Transcriptase

Notes Before use, mix each component and spin in a microcentrifuge.

It is prudent to include a No-RT Control reaction for each RNA sample by omitting the AffinityScript RT/ RNase Block enzyme mixture. This control verifies that signal detected in the subsequent QPCR is not due to genomic DNA contamination.

1. Prepare the first-strand cDNA synthesis reaction in a microcentrifuge tube by adding the following components *in order*:

RNase-free H₂O to a total volume of 20 µl
10.0 µl of first strand master mix (2×)
3.0 µl of oligo(dT) primer OR random primers (0.1 µg/µl)
1.0 µl of AffinityScript RT/ RNase Block enzyme mixture
X µl of RNA (0.3 pg–3 µg total RNA)

Note The optimum primer type [oligo(dT) or random primer] varies for different targets and should be determined empirically. See Primer Selection in PreProtocol Considerations for more information.

2. Incubate the reaction at 25°C for 5 minutes to allow primer annealing.
3. Incubate the reaction at 42°C for 15 minutes to allow cDNA synthesis.

Notes This protocol has been extensively tested and is ideal for most targets up to 12 kb. Increasing the incubation time to 45 minutes at 42°C or raising the incubation temperature from 42°C to 55°C may increase cDNA yield for longer or secondary structure-rich targets, respectively.

When using oligo(dT) to prime cDNA synthesis at an incubation temperature of 55°C, incubate the reactions at 42°C for 5 minutes before transferring to 55°C for first-strand synthesis.

4. Incubate the reactions at 95°C for 5 minutes to terminate the cDNA synthesis reaction.
5. Place the completed first-strand cDNA synthesis reactions on ice for immediate use in QPCR. For long-term storage, place the reactions at -20°C.

Real-Time PCR Quantification of First-Strand cDNA

QPCR Reagent Systems

cDNA targets may be quantified using any suitable QPCR reagent system. The AffinityScript QPCR cDNA synthesis kit is specifically qualified with Stratagene QPCR master mix kits. See Table 1 in the *Introduction* for a summary of the QPCR kits available to support different applications and detection methods.

Guidelines for cDNA Amounts for QPCR Analysis

The amount of the cDNA synthesis reaction that should be used for analysis depends on the target abundance, the QPCR reagent system used, and the amount of input RNA used. High amounts of reverse transcriptase or of input RNA originating from the cDNA synthesis reaction may inhibit the subsequent QPCR reaction.

For the Stratagene Brilliant II QPCR master mixes or Brilliant II Fast QPCR master mixes, up to 10% of the first strand synthesis reaction, containing up to 100 ng input RNA, may be added to a 25- μ l QPCR reaction. For example, if the cDNA synthesis reaction was performed with 1 μ g input RNA, up to 10% of the reaction may be added to a 25- μ l QPCR reaction. If the cDNA synthesis reaction was performed with 2 μ g input RNA, up to 5% of the reaction may be added to a 25- μ l QPCR reaction.

TROUBLESHOOTING

Observation	Suggestion
No or low yield of first-strand cDNA	<p>Verify the integrity of the input RNA by denaturing agarose gel electrophoresis to ensure it is not degraded.</p> <p>Optimize the reaction using Stratagene QPCR Reference Total RNA, which can then also be used as a calibrator for subsequent experiments.</p> <p>Prepare a new RNA sample. Use Stratagene RNA isolation kits to isolate intact total RNA or mRNA.</p> <p>Isolate the RNA in the presence of a ribonuclease inhibitor, and ensure that all RT-PCR reagents and labware are free of RNases.</p> <p>Inhibitors of reverse transcription (SDS, EDTA, guanidinium chloride, formamide, Na₂PO₄, or spermidine) may be present in the RNA sample. Reduce the volume of the input RNA or remove RT inhibitors with an additional 70% (v/v) ethanol wash following ethanol precipitation.</p> <p>Increase the length of the 42°C cDNA synthesis reaction to 45 minutes for longer RNA targets.</p> <p>Increase the incubation temperature from 42°C to 55°C for secondary structure-rich targets.</p> <p>Increase the concentration of the template RNA.</p> <p>Try switching the cDNA primer composition [oligo(dT) vs. random primers]. For challenging targets, a mixture of the two primer types may also be tested.</p> <p>When using oligo(dT) to prime cDNA synthesis at an incubation temperature of 55°C, incubate the reactions at 42°C for 5 minutes before transferring to 55°C for first-strand synthesis.</p>
No or low yield of amplification product in QPCR	<p>See the discussion under No or low yield of the first-strand cDNA for suggestions related to insufficient first strand synthesis.</p> <p>Add more cDNA synthesis product to the PCR. (Typically up to 10% of the cDNA synthesis reaction may be added to a 25-μl QPCR reaction. See Guidelines for cDNA Amounts for QPCR Analysis for additional considerations.)</p> <p>Optimize the QPCR primer concentration, annealing temperature, and/or extension time, varying each individually and in increments. Stratagene QPCR Reference Total RNAs may be used for this step to conserve experimental samples.</p> <p>Re-examine the QPCR primer design. Make sure primers are not self-complementary or complementary to each other. Verify that the primers are designed to be complementary to the appropriate strands. Try using longer primers.</p> <p>Increase the number of thermal cycles.</p> <p>To maintain adequate quality of dNTPs used in the PCR step, keep nucleotides frozen in aliquots, thaw them quickly, and keep them on ice once thawed; avoid multiple freeze-thaw cycles.</p>
Size of the amplification product is greater than expected	<p>The RNA preparation may be contaminated with genomic DNA. Test for the presence of contaminating DNA by performing RT-PCR in the absence of AffinityScript RT. If DNA contamination is confirmed, treat the RNA preparation with RNase-free DNase I. Alternatively, redesign the PCR primers to anneal to sequences in the exon-exon boundary of the target gene.</p>

REFERENCES

1. Bustin, S. A. and Nolan, T. (2004) *J Biomol Tech* 15(3):155-66.
2. Stahlberg, A., Hakansson, J., Xian, X., Semb, H. and Kubista, M. (2004) *Clin Chem* 50(3):509-15. Epub 2004 Jan 15.

ENDNOTES

SYBR® is a registered trademark of Molecular Probes, Inc.

MSDS INFORMATION

The Material Safety Data Sheet (MSDS) information for Stratagene products is provided on the web at <http://www.stratagene.com/MSDS/>. Simply enter the catalog number to retrieve any associated MSDS's in a print-ready format. MSDS documents are not included with product shipments.